

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 August 2001 (30.08.2001)

PCT

(10) International Publication Number
WO 01/62947 A1

(51) International Patent Classification⁷: **C12P 7/06**, C12N
9/16, C12C 11/00, C12G 3/02, 1/022

(21) International Application Number: PCT/DK01/00125

(22) International Filing Date: 22 February 2001 (22.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2000 00281 23 February 2000 (23.02.2000) DK

(71) Applicant: **NOVOZYMES A/S** [DK/DK]; Krogshøjvej
36, DK-2880 Bagsværd (DK).

(72) Inventors: **VEIT, Chris**; 1717 Chestnut Hill Road, Wake
Forest, NC 27587 (US). **FELBY, Claus**; Kiselvej 7E 2th,
DK-2730 Herlev (DK). **PECKOUS, Larry, W.**; 12201
Staunton Court, Raleigh, NC 27613 (US). **OLSEN, Hans,
Sejr**; Højbjerg Vang 7, DK-2840 Holte (DK).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IL,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **FERMENTATION WITH A PHYTASE**



(57) Abstract: The present invention relates to an improved fermentation process wherein phytic acid-containing material is fer-
mented in the presence of a phytase, e.g. in fermentation for the production of ethanol.

WO 01/62947 A1

FERMENTATION WITH A PHYTASE**FIELD OF THE INVENTION**

The present invention relates to a process of fermenting
5 phytic acid-containing materials; a process of alcohol and
other fermented compounds production, in particular ethanol
production; the use of phytase activity for saccharification
and/or fermentation; and a composition suitable for ethanol
production.

10 BACKGROUND OF THE INVENTION

Fermentation processes are used for making a vast number of
products of big commercial interest. Fermentation is used in
industry to produce simple compounds such as alcohols (in par-
ticular ethanol); acids, such as citric acid, itaconic acid,
15 lactic acid, gluconic acid; ketones; amino acids, such as glu-
tamic acid, but also more complex compounds such as antibiot-
ics, such as penicillin, tetracyclin; enzymes; vitamins, such
as riboflavin, B₁₂, beta-carotene; hormones, which are difficult
to produce synthetically. Also in the brewing (beer and wine
20 industry), dairy, leather, tobacco industries fermentation
processes are used.

Thus, there is a need for further improvement of fermenta-
tion processes and for processes including a fermentation step.
Accordingly, the object of the invention is to provide an im-
25 proved method at least including a fermenting step.

BRIEF DESCRIPTION OF THE DRAWING

Fig. 1 shows schematically an ethanol production process of the
invention.

Fig. 2 shows the results on fermentation of liquefied whole
30 corn mash using AMG and AMG+phytase. CO₂ loss vs. time

Fig. 3 shows the phytin level as g/kg dry matter for the corn
substrate and following fermentation with AMG and AMG + phy-
tase.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a process of producing a fermentation product, for instance the ones mentioned in the "Background of the Invention"-section, in particular ethanol, but also beverages, such as beer or wine are contemplated, wherein the fermentation is carried out in the presence of phytase activity. It is to be understood that a carbohydrate source, such as glucose, dextrose, maltose or the like, need to be present during fermentation for the fermenting organism to be able to ferment. The carbohydrate source may be supplied by direct addition of e.g., glucose, or may be supplied as a product of, e.g., (pre-)saccharification step, as will be described further below.

Ethanol production

The process of the invention may in one embodiment be an ethanol process comprising the below steps, wherein phytase activity is added during pre-saccharification and/or fermentation. It is to be understood that the phytase according to the invention may be added during the propagation of yeast cells and/or later on during the actual fermentation. Beverage production, such as beer or wine production is equally contemplated.

Alcohol production, in particular ethanol production, from whole grain can be separated into 4 main steps

- Milling
- Liquefaction
- Saccharification
- Fermentation

Milling

In one embodiment the (whole) grain is milled in order to open up the structure and allowing for further processing. Two processes are preferred according to the invention: wet and dry milling. Preferred for ethanol production is dry milling where the whole kernel is milled and used in the remaining part of the process. Wet milling may also be used and gives a good

separation of germ and meal (starch granules and protein) and is with a few exceptions applied at locations where there is a parallel production of syrups. Both dry and wet milling is well known in the art of, e.g., ethanol production.

5 Liquefaction

In an embodiment of the liquefaction step of the invention, milled gelatinized whole grain raw material is broken down (hydrolyzed) into maltodextrins (dextrins) mostly of a DE higher than 4. The hydrolysis may be carried out by acid treatment or
10 enzymatically by alpha-amylase treatment, in particular with a *Bacillus* alpha-amylase as will be described further below. Acid hydrolysis is used on a limited basis. The raw material is in one embodiment of the invention milled (whole) grain. However, a side stream from starch processing may also be used.

15 In an embodiment of the invention enzymatic liquefaction is carried out as a three-step hot slurry process. The slurry is heated to between 60-95°C, preferably 80-85°C (in the Slurry Tank - see Fig. 1), and the enzyme(s) is(are) added to initiate liquefaction (thinning). Then the slurry is jet-cooked at a
20 temperature between 95-140°C, preferably 105-125°C to complete gelatinization of the slurry. Then the slurry is cooled to 60-95°C and more enzyme(s) is(are) added to finalize hydrolysis (secondary liquefaction). The liquefaction process is carried out at pH 4.5-6.5, in particular at a pH between 5 and 6.
25 Milled and liquefied whole grains are known as mash.

Saccharification

To produce low molecular sugars DP_{1-3} that can be metabolized by yeast, the maltodextrin from the liquefaction must be further hydrolyzed. The hydrolysis is typically done enzymatically
30 by glucoamylases, alternatively alpha-glucosidases or acid alpha-amylases can be used. A full saccharification step may last up to 72 hours, however, it is common only to do a pre-saccharification of typically 40-90 minutes and then complete
35 saccharification during fermentation (SSF). Saccharification is

typically carried out at temperatures from 30-65°C, typically around 60°C, and at pH 4.5.

Fermentation

5 Yeast typically from *Saccharomyces* spp. is added to the mash and the fermentation is ongoing for 24-96 hours, such as typically 35-60 hours. The temperature is between 26-34°C, in particular about 32°C, and the pH is from pH 3-6, preferably around pH 4-5.

10 Note that the most widely used process is a simultaneous saccharification and fermentation (SSF) process where there is no holding stage for the saccharification, meaning that fermenting organism, such as the yeast, and enzyme(s) is(are) added together. When doing SSF it is common to introduce a pre-
15 saccharification step at a temperature above 50°C, just prior to the fermentation.

Distillation

Optionally following the fermentation, the mash may be dis-
20 tilled to extract the, for instance, ethanol.

In the case where the end product is ethanol, obtained according to the process of the invention, it may be used as, e.g., fuel ethanol; drinking ethanol, i.e., potable neutral spirits; or industrial ethanol.

25

By-products

Left over from the fermentation is the grain, which is typically used for animal feed either in liquid or dried form.

Further details on how to carry out liquefaction, sacchari-
30 fication, fermentation, distillation, and recovering of ethanol are well known to the skilled person.

According to the process of the invention the saccharification and fermentation may be carried out simultaneously or separately.

35 In the first aspect the invention relates to a process, wherein phytic acid-containing material is fermented in the

presence of a phytase. The addition of a phytase during the fermentation (or e.g., a combined or simultaneous fermentation and saccharification step, as will be described further below) results in a number of advantages. It is to be understood that the phytase according to the invention may be added during the propagation of yeast cells and/or later on during the actual fermentation. For instance, the addition of phytase results in that more free minerals, e.g., Ca^{2+} , are made available to the fermenting organism(s), in particular yeast. Phytic acid interferes with minerals such as especially calcium, magnesium, zinc and iron, in a manner that reduce the stability of a number of alpha-amylases, in particular *Bacillus* (e.g., BAN or *Bacillus licheniformis* alpha-amylases) or *Aspergillus* (e.g., the *Aspergillus oryzae* alpha-amylase sold as FUNGAMYL™ from Novozymes A/S) alpha-amylases. Therefore, a higher fermentation rate is a result of the hydrolysis of phytin making accessible minerals essential to the yeast metabolism.

In preferred embodiment the addition of phytase is made in combination with a "Carbohydrate-source generating enzyme". The term "carbohydrate-source generating enzyme" includes gluco-amylases (being a glucose generator), and beta-amylases and maltogenic amylases (being maltose generators).

The carbohydrate-source generating enzymes are capable of providing energy to the fermenting microorganism(s) in question.

Further, the availability of more free phosphorus minerals and vitamins - as a result of phytin to inositol conversion - improves the yeast growth and viability during fermentation and thus increases the fermentation and ethanol yields.

Further, the protein availability is increased.

The by-product of an ethanol process may be used as feed (Distillers Dried Grain) and has an improved nutritional value in comparison to a corresponding process where a phytase has not been added.

In one embodiment the invention relates to a process for the production of ethanol, comprising the steps of:

(a) milling whole grains,

(b) liquefying the product of step (a), in the presence of an alpha-amylase,

(c) saccharifying the liquefied material obtained in step (b) in the presence of a phytase,

5 (d) fermenting the saccharified material obtained in step (c) using a micro-organism; and optionally

(e) distilling of the fermented and saccharified material obtained in step (d).

10 In a preferred embodiment the whole grains in step a) are dry milled, for instance in a hammer mill.

In an embodiment the DS% (dry solid percentage) in the slurry tank (containing milled whole grains) is in the range from 1-60%, in particular 10-50%, such as 20-40%, such as 25-35%.

15 In a preferred embodiment of the invention the liquefaction step comprising the following sub-steps:

b1) the hot slurry is heated to between 60-95°C, preferably 80-85°C, and at least an alpha-amylase is added;

20 b2) the slurry is jet-cooked at a temperature between 95-140°C, preferably 105-125°C to complete gelanitization of the slurry;

b3) the slurry is cooled to 60-95°C and more alpha-amylase is added to finalize hydrolysis.

25 The liquefaction process is in an embodiment carried out at pH 4.5-6.5, in particular at a pH between 5 and 6.

Steps (c) and (d) may be carried out either simultaneously or separately/sequential. Further, after step (e) an optional ethanol recovery step may be added.

30 Materials Used in Processes of the Invention

Phytic acid-containing material

According to the above described processes of the invention the raw materials contain phytic acids and fermentable sugars or constituents, which can be converted into sugars. This in-
35 clude starch-containing raw materials, such as tubers, roots, whole grains, corns, cobs, wheat, barley, rye, milo or cereals, sugar-containing raw materials, such as molasses, fruit materi-

als, sugar, cane or sugar beet, potatoes, cellulose-containing materials, such as wood or plant residues.

According to the invention the phytic acid containing material may be the side stream from starch processing, in particular liquefied starch with a DE of 6-20, in particular between
5 8-10.

Microorganisms for fermentation

The microorganism may be a fungal organism, such as yeast or bacteria. Examples of filamentous fungi include strains of
10 *Penicillium* sp. Preferred organisms for ethanol production is yeasts. Preferred yeast according to the invention is baker's yeast, also known as *Saccharomyces cerevisiae*. The yeast may according to the invention preferably be added before starting the actual fermentation (i.e., during the propagation phase).
15 The yeast cells may be added in amounts of 10^5 to 10^{12} , preferably from 10^7 to 10^{10} , especially 5×10^7 viable yeast count per ml of fermentation broth. During the ethanol producing phase the yeast cell count should preferably be in the range from 10^7 to 10^{10} , especially around 2×10^8 . Example 1 shows a fermentation
20 process of the invention where the yeast is not stressed (yeast count of about 10^{10} cells per ml). Even under such conditions the addition of phytase is shown to improved the fermentation process. Further guidance in respect of using yeast for fermentation can be found in, e.g., "The alcohol Textbook" (Editors
25 K. Jacques, T.P. Lyons and D.R.Kelsall, Nottingham University Press, United Kingdom 1999), which is hereby incorporated by reference.

ENZYMES

30 Phytase

The phytase used according to the invention may be any enzyme capable of effecting the liberation of inorganic phosphate from phytic acid (myo-inositol hexakisphosphate) or from any salt thereof (phytates). Phytases can be classified according
35 to their specificity in the initial hydrolysis step, viz. according to which phosphate-ester group is hydrolyzed first. The

phytase to be used in the invention may have any specificity, e.g., be a 3-phytase (EC 3.1.3.8), a 6-phytase (EC 3.1.3.26) or a 5-phytase (no EC number).

In a preferred embodiment the phytase has a temperature optimum in the range from 25-70°C, preferably 28-50°C, especially 30-40°C. This is advantageous when the phytase is added during fermentation.

In another preferred embodiment the phytase has a temperature optimum above 50°C, such as in the range from 50-70°C. This is advantageous when the phytase is added during (pre-)saccharification. The dosage of the phytase may be in the range 5.000-250.000 FYT/g DS, particularly 10.000-100.000 FYT/g DS. A preferred suitable dosage of the phytase is in the range from 0.005-25 FYT/g DS, preferably 0.01-10 FYT/g, such as 0.1-1 FYT/g DS.

Here, the phytase activity is determined FYT units, one FYT being the amount of enzyme that liberates 1 micromole inorganic ortho-phosphate per min. under the following conditions: pH 5.5; temperature 37°C; substrate: sodium phytate ($C_6H_6O_{24}P_6Na_{12}$) at a concentration of 0.0050 mole/l.

The phytase may be derived from plants or microorganisms, such as bacteria or fungi, e.g., yeast or filamentous fungi.

The plant phytase may be from wheat-bran, maize, soy bean or lily pollen. Suitable plant phytases are described in Thomlinson et al, Biochemistry, 1 (1962), 166-171; Barrientos et al, Plant. Physiol., 106 (1994), 1489-1495; WO 98/05785; WO 98/20139.

A bacterial phytase may be from genus *Bacillus*, *Pseudomonas* or *Escherichia*, specifically the species *B. subtilis* or *E. coli*. Suitable bacterial phytases are described in Paver and Jagannathan, 1982, Journal of Bacteriology 151:1102-1108; Cosgrove, 1970, Australian Journal of Biological Sciences 23:1207-1220; Greiner et al, Arch. Biochem. Biophys., 303, 107-113, 1993; WO 98/06856; WO 97/33976; WO 97/48812.

A yeast phytase or myo-inositol monophosphatase may be derived from genus *Saccharomyces* or *Schwanniomyces*, specifically species *Saccharomyces cerevisiae* or *Schwanniomyces occiden-*

talis. The former enzyme has been described as a Suitable yeast phytases are described in Nayini et al, 1984, Lebensmittel Wissenschaft und Technologie 17:24-26; Wodzinski et al, Adv. Appl. Microbiol., 42, 263-303; AU-A-24840/95;

5 Phytases from filamentous fungi may be derived from the fungal phylum of *Ascomycota* (ascomycetes) or the phylum *Basidiomycota*, e.g., the genus *Aspergillus*, *Thermomyces* (also called *Humicola*), *Myceliophthora*, *Manascus*, *Penicillium*, *Peniophora*, *Agrocybe*, *Paxillus*, or *Trametes*, specifically the species *Aspergillus terreus*, *Aspergillus niger*, *Aspergillus niger*
10 var. *awamori*, *Aspergillus ficuum*, *Aspergillus fumigatus*, *Aspergillus oryzae*, *T. lanuginosus* (also known as *H. lanuginosa*), *Myceliophthora thermophila*, *Peniophora lycii*, *Agrocybe pediades*, *Manascus anka*, *Paxillus involtus*, or *Trametes pubescens*. Suitable fungal phytases are described in Yamada et al., 1986, Agric. Biol. Chem. 322:1275-1282; Piddington et al., 1993, Gene 133:55-62; EP 684,313; EP 0 420 358; EP 0 684 313; WO 98/28408; WO 98/28409; JP 7-67635; WO 98/44125; WO 97/38096; WO 98/13480.

20 Modified phytases or phytase variants are obtainable by methods known in the art, in particular by the methods disclosed in EP 897010; EP 897985; WO 99/49022; WO 99/48330.

 Commercially available phytases contemplated according to the invention include BIO-FEED PHYTASE™, PHYTASE NOVO™ CT or L
25 (all from Novozymes), or NATUPHOS™ NG 5000 (from DSM).

Alpha-amylase

 The liquefaction step may be performed in the presence of an alpha-amylase derived from a microorganism or a plant. Preferred alpha-amylases are of fungal or bacterial origin. *Bacillus* alpha-amylases (often referred to as "Termamyl-like alpha-amylases"), variant and hybrids thereof, are specifically contemplated according to the invention. Well-known Termamyl-like alpha-amylases include alpha-amylase derived from a strain of *B. licheniformis* (commercially available as Termamyl™), *B. amyloliquefaciens*, and *B. stearothermophilus* alpha-amylase (BSG).
35 Other Termamyl-like alpha-amylases include alpha-amylase derived

from a strain of the *Bacillus* sp. NCIB 12289, NCIB 12512, NCIB 12513 or DSM 9375, all of which are described in detail in WO 95/26397, and the alpha-amylase described by Tsukamoto et al., Biochemical and Biophysical Research Communications, 151 (1988), pp. 25-31. In the context of the present invention a Termamyl-like alpha-amylase is an alpha-amylase as defined in WO 99/19467 on page 3, line 18 to page 6, line 27. Contemplated variants and hybrids are described in WO 96/23874, WO 97/41213, and WO 99/19467. Contemplated alpha-amylase derived from a strain of *Aspergillus* includes *Aspergillus oryzae* and *Aspergillus niger* alpha-amylases. Commercial alpha-amylase products and products containing alpha-amylases include TERMAMYL™ SC, FUNGAMYL™, LIQUOZYME™ and SAN™ SUPER.

Fungal alpha-amylases may be added in an amount of 0.001-1.0 AFAU/g DS, preferably from 0.002-0.5 AFAU/g DS, preferably 0.02-0.1 AFAU/g DS.

Bacillus alpha-amylases may be added in effective amounts well known to the person skilled in the art.

20 Glucoamylase

The saccharification step (c) or a combined saccharification and fermentation step (SSF step) may be carried out in the presence of a glucoamylase derived from a microorganism or a plant. Preferred is glucoamylase of fungal or bacterial origin selected from the group consisting of *Aspergillus* glucoamylases, in particular *A. niger* G1 or G2 glucoamylase (Boel et al. (1984), EMBO J. 3 (5), p. 1097-1102), or variants thereof, such as disclosed in WO 92/00381 and WO 00/04136; the *A. awamori* glucoamylase (WO 84/02921), *A. oryzae* (Agric. Biol. Chem. (1991), 55 (4), p. 941-949), or variants or fragments thereof.

Other contemplated *Aspergillus* glucoamylase variants include variants to enhance the thermal stability: G137A and G139A (Chen et al. (1996), Prot. Engng. 9, 499-505); D257E and D293E/Q (Chen et al. (1995), Prot. Engng. 8, 575-582); N182 (Chen et al. (1994), Biochem. J. 301, 275-281); disulphide bonds, A246C (Fierobe et al. (1996), Biochemistry, 35, 8698-

8704; and introduction of Pro residues in position A435 and S436 (Li et al. (1997), *Protein Engng.* 10, 1199-1204. Furthermore Clark Ford presented a paper on Oct 17, 1997, ENZYME ENGINEERING 14, Beijing/China Oct 12-17, 97, Abstract number: 5 Abstract book p. 0-61. The abstract suggests mutations in positions G137A, N20C/A27C, and S30P in an *Aspergillus awamori* glucoamylase to improve the thermal stability. Other glucoamylases include *Talaromyces* glucoamylases, in particular derived from *Talaromyces emersonii* (WO 99/28448), *Talaromyces leycettanus* 10 (US patent no. Re. 32,153), *Talaromyces duponti*, *Talaromyces thermophilus* (US patent no. 4,587,215). Bacterial glucoamylases contemplated include glucoamylases from the genus *Clostridium*, in particular *C. thermoamylolyticum* (EP 135,138), and *C. thermohydrosulfuricum* (WO 86/01831).

15 Glucoamylases may in an embodiment be added in an amount of 0.02-2 AGU/g DS, preferably 0.1-1 AGU/g DS, such as 0.2 AGU/g DS

The ratio between acidic fungal alpha-amylase activity (AFAU) per glucoamylase activity (AGU) (AFAU per AGU) may in 20 one embodiment be at least 0.1, in particular at least 0.16, such as in the range from 0.12 to 0.30.

Commercial products include SANTM SUPER and AMGTM E (from Novozymes); OPTIDEXTM 300 (from Genencor Int.); AMIGASETM (from DSM); G-ZYMETM G900 (from Enzyme Bio-Systems). 25

Protease

Addition of protease(s) increase(s) the FAN (Free amino nitrogen) level and increase the rate of metabolism of the yeast and further gives higher fermentation efficiency.

30 Suitable proteases include fungal and bacterial proteases. Preferred proteases are acidic proteases, i.e., proteases characterized by the ability to hydrolyze proteins under acidic conditions below pH 7.

Suitable acid fungal proteases include fungal proteases derived from *Aspergillus*, *Mucor*, *Rhizopus*, *Candida*, *Coriolus*, *Endothia*, *Enthomophtra*, *Irpex*, *Penicillium*, *Sclerotium* and *Torulopsis*. Especially contemplated are proteases derived from As- 35

Aspergillus niger (see, e.g., Koaze et al., (1964), *Agr. Biol. Chem. Japan*, 28, 216), *Aspergillus saitoi* (see, e.g., Yoshida, (1954) *J. Agr. Chem. Soc. Japan*, 28, 66), *Aspergillus awamori* (Hayashida et al., (1977) *Agric. Biol. Chem.*, 42(5), 927-933, 5 *Aspergillus aculeatus* (WO 95/02044), or *Aspergillus oryzae*; and acidic proteases from *Mucor pusillus* or *Mucor miehei*.

Bacterial proteases, which are not acidic proteases, include the commercially available products Alcalase® and Neutrase® (available from Novozymes A/S).

10 Protease(s) may in one embodiment be added in an amount of 10^{-7} to 10^{-5} gram active protease protein/g DS, in particular 10^{-7} to 5×10^{-6} gram active protease protein/g DS

Additional enzyme

15 One or more additional enzymes may also be incorporated. Additional enzymes include pullulanases.

In a final aspect the invention relates to the use of phytase, in particular the phytases mentioned above for saccharification and/or fermentation (SSF) and for use for ethanol 20 production.

Finally the invention related to a composition comprising a phytase and at least one carbohydrate-source generating enzyme (as defined above), in particular a glucoamylase, such as an *Aspergillus niger* and/or *Talaromuces emersonii* glucoamylase. 25 The composition may further comprise a protease, in particular an acid protease, such as an acid fungal protease.

MATERIALS AND METHODS

Materials:

30 Phytase: derived from *Peniophora lycii* (LN 0 1305 dept. 550) E-1999-0082) (available from Novozymes)

Glucoamylase: *A. niger* glucoamylase (available as AMG E (1999-SE-0025) from Novozymes)

Alpha-amylase: BSG (*B. stearothermophilus* alpha-amylase which 35 is available from Novozymes as TERMAMYL™ SC)

Mash: Liquefied whole corn mash prepared by a hot slurry process and Termamyl SC. The mash had a DE of about 17 and a dry substance of 28%.

Yeast: Dry yeast (*Saccharomyces cerevisiae*)

5

Determination of Alpha-Amylase Activity (KNU)

1. Phadebas assay

Alpha-amylase activity is determined by a method employing Phadebas® tablets as substrate. Phadebas tablets (Phadebas® Amylase Test, supplied by Pharmacia Diagnostic) contain a cross-linked insoluble blue-colored starch polymer, which has been mixed with bovine serum albumin and a buffer substance and tableted.

For every single measurement one tablet is suspended in a tube containing 5 ml 50 mM Britton-Robinson buffer (50 mM acetic acid, 50 mM phosphoric acid, 50 mM boric acid, 0.1 mM CaCl₂, pH adjusted to the value of interest with NaOH). The test is performed in a water bath at the temperature of interest. The alpha-amylase to be tested is diluted in x ml of 50 mM Britton-Robinson buffer. 1 ml of this alpha-amylase solution is added to the 5 ml 50 mM Britton-Robinson buffer. The starch is hydrolyzed by the alpha-amylase giving soluble blue fragments. The absorbance of the resulting blue solution, measured spectrophotometrically at 620 nm, is a function of the alpha-amylase activity.

It is important that the measured 620 nm absorbance after 10 or 15 minutes of incubation (testing time) is in the range of 0.2 to 2.0 absorbance units at 620 nm. In this absorbance range there is linearity between activity and absorbance (Lambert-Beer law). The dilution of the enzyme must therefore be adjusted to fit this criterion. Under a specified set of conditions (temp., pH, reaction time, buffer conditions) 1 mg of a given alpha-amylase will hydrolyze a certain amount of substrate and a blue colour will be produced. The colour intensity is measured at 620 nm. The measured absorbance is directly proportional to the spe-

cific activity (activity/mg of pure alpha-amylase protein) of the alpha-amylase in question under the given set of conditions.

2. Alternative method

Alpha-amylase activity is determined by a method employing the PNP-G7 substrate. PNP-G7 which is a abbreviation for p-nitrophenyl-alpha,D-maltoheptaoside is a blocked oligosaccharide which can be cleaved by an endo-amylase. Following the cleavage, the alpha-Glucosidase included in the kit digest the substrate to liberate a free PNP molecule which has a yellow colour and thus can be measured by visible spectrophotometry at $\lambda=405\text{nm}$. (400-420 nm.). Kits containing PNP-G7 substrate and alpha-Glucosidase is manufactured by Boehringer-Mannheim (cat. No. 1054635).

To prepare the substrate one bottle of substrate (BM 1442309) is added to 5 ml buffer (BM1442309). To prepare the alpha-Glucosidase one bottle of alpha-Glucosidase (BM 1462309) is added to 45 ml buffer (BM1442309). The working solution is made by mixing 5 ml alpha-Glucosidase solution with 0.5 ml substrate.

The assay is performed by transforming 20 micro l enzyme solution to a 96 well microtitre plate and incubating at 25°C. 200 micro l working solution, 25°C is added. The solution is mixed and pre-incubated 1 minute and absorption is measured every 15 sec. over 3 minutes at OD 405 nm.

The slope of the time dependent absorption-curve is directly proportional to the specific activity (activity per mg enzyme) of the alpha-amylase in question under the given set of conditions.

Determination of Acid Amylolytic Activity (FAU)

One Fungal Alpha-Amylase Unit (1 FAU) is defined as the amount of enzyme, which breaks down 5.26 g starch (Merck Amylum solubile Erg. B.6, Batch 9947275) per hour at Novo Nordisk's standard method for determination of alpha-amylase based upon the following standard conditions:

Substrate Soluble starch
Temperature 37°C

pH. 4.7

Reaction time 7-20 minutes

A detailed description of Novo Nordisk's method is available on request.

5

Determination of acid alpha-amylase activity (AFAU)

Acid alpha-amylase activity is measured in AFAU (Acid Fungal Alpha-amylase Units), which are determined relative to an enzyme standard.

10 The standard used is AMG 300 L (wild type *A. niger* G1 AMG sold by Novo Nordisk). The neutral alpha-amylase in this AMG falls after storage at room temperature for 3 weeks from approx. 1 FAU/mL to below 0.05 FAU/mL.

15 The acid alpha-amylase activity in this AMG standard is determined in accordance with AF 9 1/3 (available from Novo method for the determination of fungal alpha-amylase). In this method, 1 AFAU is defined as the amount of enzyme, which degrades 5.260 mg starch dry matter per hour under standard conditions.

20 Iodine forms a blue complex with starch but not with its degradation products. The intensity of colour is therefore directly proportional to the concentration of starch. Amylase activity is determined using reverse colorimetry as a reduction in the concentration of starch under specified analytic conditions.
25

Alpha-amylase

Starch + Iodine → Dextrins + Oligosaccharides

40°C, pH 2.5

Blue/violet t=23 sec. Decolouration

30

Standard conditions/reaction conditions: (per minute)

Substrate: starch, approx. 0.17 g/L

Buffer: Citrate, approx. 0.03 M

Iodine (I₂): 0.03 g/L

35 CaCl₂: 1.85 mM

pH: 2.50 ± 0.05

Incubation temperature: 40°C

Reaction\time: 23 seconds
Wavelength: Lambda=590nm
Enzyme concentration: 0.025 AFAU/mL
Enzyme working range: 0.01-0.04 AFAU/mL

- 5 Further details can be found in EB-SM-0259.02/01 available on request from Novozymes, and hereby incorporated by reference.

Determination Of Glucoamylase Activity (AGU)

- 10 The Novo Amyloglucosidase Unit (AGU) is defined as the amount of enzyme, which hydrolyzes 1 micromole maltose per minute at 37°C and pH 4.3. A detailed description of the analytical method (AEL-SM-0131) is available on request from Novo Nordisk.

- 15 The activity is determined as AGU/ml by a method modified after (AEL-SM-0131, available on request from Novo Nordisk) using the Glucose GOD-Perid kit from Boehringer Mannheim, 124036. Standard: AMG-standard, batch 7-1195, 195 AGU/ml. 375 microL substrate (1% maltose in 50 mM Sodium acetate, pH 4.3) is incubated 5 minutes at 37°C. 25 microL enzyme diluted in sodium acetate is added. The reaction is stopped after 10 minutes by adding 100 microL 0.25 M NaOH. 20 microL is transferred to a 96 well microtitre plate and 200 microL GOD-Perid solution (124036, Boehringer Mannheim) is added. After 30 minutes at room temperature, the absorbance is measured at 650 nm and the activity calculated in AGU/ml from the AMG-standard.
- 20
25

EXAMPLES

- 30 Example 1

Fermentation in the presence of phytase

- 250 g of mash was filled into a 500 mL blue cap bottle. The pH of the mash was adjusted to 4.5. Before fermentation a pre-saccharification step was carried out by adding the saccharification enzymes and placing the bottles in a water bath at 60°C for 70 minutes. The bottle was cooled in a water bath for 40
- 35

minutes to 30°C and dry yeast was added at a dosage of 0.8 g/bottle (in order to reach 30°C within 40 minutes ice is added to the water bath). The bottle is closed by a yeast-lock filled with concentrated H₂SO₄. The fermentation was continued for 96
 5 hours and the fermentation rate was monitored by weighing the bottle at regular intervals for measuring CO₂ loss.

Determination of phytin content

The phytin content was measured for all treatments and the
 10 substrate. The phytin content was measured by The Danish Institute of Agricultural sciences, Tjele, Denmark, according to the method described in: Brooks, J.R. and C.V. Morr. 1984. Phosphorus and phytate content of soybean protein components. J. Agric. Food Chem. 32: 872-874.

15

Treatments

Three different treatments were applied (all enzyme dosages are based on dry matter):

0.2 AGU/g AMG E

20 0.2 AGU/g AMG E + 1.0 FYT/g phytase

1.0 FYT/g phytase

Each treatment was duplicated 4 times.

The results of the fermentations are shown in Fig. 2 and
 25 Table 1. The positive effect of the phytase is significant at a level of P=0.05. The effect of the phytase treatment on the level of phytin is shown in Fig. 3. The phytase efficiently hydrolyses the phytin below the detection level.

Hours	CO ₂ (g)		
	0.2 AGU/g	0.2 AGU/g + 1 FYT/g	1 FYT/g
CO ₂ loss 96 hr.	23.0	23.9	5.3
% of theoretical yield at 96 hours	88.0	91.1	20.2

Table 1. Fermentation of whole corn mash. CO₂ loss after 96 hour fermentation using AMG E and phytase.

CLAIMS

1. A process including a fermentation step, wherein phytic acid-containing material is fermented in the presence of a phytase.

5 2. The process of claim 1, wherein the phytase is derived from a strain of *Peniophra lycii*, *Aspergillus oryzae*, *Aspergillus ficumum*, or *Aspergillus niger*.

3. The process of claim 1-2, wherein the phytase has a temperature optimum in the range 20-70°C, in particular above 50°C.

10 4. The process of claims 1-3, wherein the phytic acid-containing materials are fermented with a micro organism capable of fermenting sugars or converted sugars.

5. The process of claim 4, wherein the micro-organism is a yeast, such as *Saccharomyces cerevisae*.

15 6. The process of claims 1-5, wherein the phytic acid-containing material derived from whole grains or a side stream from starch processing, in particular liquefied starch with a DE of 8-10.

20 7. The process of claim 1-6, wherein the fermentation is carried out in the presence of a carbohydrate-generating enzyme, such as a glucoamylase.

8. The process of claim 1-7, wherein the fermentation is carried out in the presence of a protease, in particular a protease selected from the group of fungal proteases, such as an acid fungal protease derived from a strain of *Aspergillus*, in particular *A. niger*.

9. The process of claim 1-7, wherein the fermentation is carried out in the presence of a protease, in particular a protease selected from the group of bacterial proteases, such as a acidic, neutral or alkaline protease, such as a protease derived from a strain of *Bacillus*, in particular Alcalase® or

30

Neutrase®.

10. The process of claim 1-9, wherein the fermentation product is ethanol.

11. A process for the production of ethanol, comprising the
5 steps of:

(a) milling whole grains,

(b) liquefying the product of step (a), in the presence of an alpha-amylase,

(c) saccharifying the liquefied material obtained in step
10 (b),

(d) fermenting the saccharified material obtained in step (c) using a micro-organism, and optionally

(e) distilling of the fermented and saccharified material obtained in step (d).

15 12. The process of claim 11, wherein a carbohydrate-source generating enzyme, such as a glucoamylase is added during saccharification.

13. The process of claims 1-12, wherein the liquefaction step comprising the following sub-steps:

20 b1) the hot slurry is heated to between 60-95°C, preferably 80-85°C, and at least an alpha-amylase is added;

b2) the slurry is jet-cooked at a temperature between 95-140°C, preferably 105-125°C to complete gelatinization of the slurry;

25 b3) the slurry is cooled to 60-95°C and more alpha-amylase is added to finalize hydrolysis.

14. The process of claims 1-13, wherein the liquefaction process is carried out at pH 4.5-6.5, in particular at a pH between 5 and 6.

15. The process of claims 1-14, wherein the phytase is added during pre-saccharification or saccharification.

16. The process of claims 1-15, wherein a phytase is added during saccharification and/or fermentation.

5 17. The process of claims 11-16, wherein steps (c) and (d) is carried out either simultaneously or separately/sequentially.

18. The process of claims 11-17, wherein an optionally ethanol recovery step is carried out after step (e).

10 19. The process of claim 11-18, wherein the milling step (a) is a dry milling step or wet milling step.

20. The process of claims 11-1, wherein the alpha-alpha-amylase is derived from a strain of the genus *Bacillus* or a strain of *Aspergillus*.

15 21. The process of claims 11-20, wherein the phytase is derived from a strain of *Peniophra lycii* or *Aspergillus oryzae*

22. The process of claims 11-21, wherein the fermentation is carried out using a micro-organism capable of fermenting sugars to ethanol.

20 23. The process of claim 22, wherein the micro-organism is a yeast, such as *Saccharomyces cerevisiae*.

24. The process of claims 11-23, wherein the phytic acid-containing material is milled whole grain or a side stream from starch processing.

25 25. The process of claims 11-24, wherein the fermentation is carried out in the presence of carbohydrate-source generating enzyme, such as a glucoamylase, and/or further a protease.

26. The process of claim 23, wherein the protease is selected from the group of fungal proteases, such as an acid fungal protease derived from a strain of *Aspergillus*, in particu-

30

lar *A. niger*.

27. The process of claim 26, wherein the protease is an acid, neutral or alkaline protease, such as a protease derived from a strain of *Bacillus*, in particular Alcalase® or Neutrase®.

28. Use of phytase for saccharification and/or fermentation.

29. Use of phytase for ethanol production.

10

30. A composition comprising a phytase and at least one carbohydrate-generating enzyme, in particular a glucoamylase.

31. A composition of claim 30, said composition further comprising a protease, in particular an acid protease, especially an acid fungal protease.




Fig. 1

2/3




Fig. 2

BEST AVAILABLE COPY

3/3




Fig. 3

BEST AVAILABLE COPY

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 01/00125

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C12P 7/06, C12N 9/16, C12C 11/00, C12G 3/02, C12G 1/022
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C12P, C12C, C12G, C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO INTERNAL, WPI DATA, BIOSIS, BIOTECHNOLOGY ABSTRACTS, CHEM. ABS DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9613600 A1 (GENECOR INTERNATIONAL, INC.), 9 May 1996 (09.05.96), claims --	1-31
X	WO 9220777 A1 (SOLVAY ENZYMES INC.), 26 November 1992 (26.11.92), claims --	11-14, 17-20, 22-27
X	US 5756714 A (ANTRIM ET AL), 26 May 1998 (26.05.98), column 2, line 10 - line 19; column 6, line 41 - line 49 --	1-3, 7, 13-14, 18-21
A	US 5902615 A (SANIEZ ET AL), 11 May 1999 (11.05.99), column 1, line 12 - line 13; column 1, line 33 - line 34; column 5, line 21 - line 25 --	1-31

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

13 June 2001

Date of mailing of the international search report

29.06.2001

Name and mailing address of the International Searching Authority
 European Patent Office P.B. 6818 Patentlaan 2
 NL-2280 HV Rijswijk
 Tel (+31-70) 340-2040, Tx 31 651 epo nl,
 Fax (+31-70) 340-3016
 E-mail: epo@epo.nl

Authorized officer

Carl-Olof Gustafsson/BS
 Telephone No. --

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 01/00125

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0321004 A1 (DORR-OLIVER INCORPORATED), 21 June 1989 (21.06.89), page 1, line 10 - line 11, claims 1-6 -- -----	1-31

INTERNATIONAL SEARCH REPORT

Information on patent family members

28/05/01

International application No.

PCT/DK 01/00125

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9613600	A1	09/05/96	AU	4013295 A	23/05/96
				CA	2203811 A	09/05/96
				EP	0788551 A	13/08/97
				FI	971781 A	25/04/97
WO	9220777	A1	26/11/92	AT	170217 T	15/09/98
				AU	661702 B	03/08/95
				AU	2155692 A	30/12/92
				BR	9206015 A	27/12/94
				CA	2102896 A	18/11/92
				CN	1045996 B	27/10/99
				CN	1069768 A	10/03/93
				CN	1231926 A	20/10/99
				DE	69226772 D,T	14/01/99
				DK	585374 T	25/05/99
				EP	0585374 A,B	09/03/94
				JP	7509118 T	12/10/95
				US	5231017 A	27/07/93
US	5756714	A	26/05/98	AU	710631 B	23/09/99
				AU	5298496 A	02/10/96
				BR	9607474 A	23/12/97
				CA	2214383 A	19/09/96
				EP	0813607 A	29/12/97
				JP	11501813 T	16/02/99
				NO	974072 A	29/10/97
				NZ	305062 A	28/10/99
				WO	9628567 A	19/09/96
US	5902615	A	11/05/99	AT	197159 T	15/11/00
				CA	2210511 A	18/01/98
				DE	69703371 D,T	03/05/01
				DK	819702 T	05/02/01
				EP	0819702 A,B	21/01/98
				ES	2152072 T	16/01/01
				FR	2751333 A,B	23/01/98
				NO	973291 A	19/01/98

INTERNATIONAL SEARCH REPORT

Information on patent family members

28/05/01

International application No.

PCT/DK 01/00125

Patent document cited in search report				Publication date		Patent family member(s)	Publication date
EP	0321004	A1	21/06/89	AT	71811	T	15/02/92
				CA	1296668	A	03/03/92
				CN	1018518	B	07/10/92
				CN	1033925	A	19/07/89
				DE	3868032	A	05/03/92
				ES	2028263	T	01/07/92
				FI	93919	B,C	15/03/95
				FI	884046	A	18/05/89
				JP	1281048	A	13/11/89
				JP	2589560	B	12/03/97
				KR	124792	B	27/11/97
				MX	26691	A	01/06/93
				MX	173082	B	31/01/94
				NL	8702735	A	16/06/89
				PH	24663	A	07/09/90
				PT	88316	A,B	01/09/88
				SU	1829959	A	23/07/93
				US	4914029	A	03/04/90
				ZA	8805074	A	29/03/89